

HIGHER ORDER CAYLEY TRANSFORMS WITH APPLICATIONS TO ATTITUDE REPRESENTATIONS

Panagiotis Tsiotras *

University of Virginia
Charlottesville, VA 22903-2442

John L. Junkins † and Hanspeter Schaub ‡

Texas A&M University
College Station, TX 77843-3141

Abstract

In this paper we generalize some previous results on attitude representations using Cayley transforms. First, we show that proper orthogonal matrices, that naturally represent rotations, can be generated by a form of “conformal” analytic mappings in the space of matrices. Using a natural parallelism between the elements of the complex plane and the real matrices, we generate higher order Cayley transforms and we discuss some of their properties. These higher order Cayley transforms are shown to parameterize proper orthogonal matrices into higher order “Rodrigues” parameters.

1. Introduction

The question of the proper choice of coordinates for describing rotations has a very long and exciting history. Starting with the work of Euler and Hamilton a series of different parameterizations were introduced by several researchers during the past hundred years. We will not delve into these results here since they can be found in any good textbook on attitude representations^{1,2}. We just mention the recent survey article by Shuster³ in the special issue in Ref. [4].

In this paper we take a slightly more abstract point of view than the previous references. Our main objective is to “unify” some of the existing results in the area of attitude representations. It is hoped that this global view will add to the current understanding of attitude representations. Our motivation stems mainly from the recent results on second order Rodrigues parameters^{5,6,7}. In particular, in Ref. [7] it was shown that these (Modified) Rodrigues parameters can be generated by a second

order Cayley transform, the same way the classical Cayley-Rodrigues parameters are generated by the Cayley transform⁸. Viewing the Cayley transform as a bilinear transformation which maps the space of skew-symmetric matrices onto the space of proper orthogonal matrices (and vice versa) one is naturally led to the notion of conformal mappings (a generalization of the bilinear transformation) from the imaginary axis onto the unit circle (and vice versa). We seek to generalize these conformal mappings to matrix spaces. Drawing on the insightful statements by Halmos⁹ we show that such an intuitive generalization is indeed possible. We are therefore able to generate the Euler parameters, the Rodrigues parameters and the Modified Rodrigues parameters as special cases of such conformal mappings. Higher order Rodrigues parameters can be easily constructed using this approach, although their relevance to applications is still to be determined. We explicitly develop the third and fourth order “Rodrigues parameters” in order to illustrate potential advantages as well as difficulties. The question of kinematics of these higher order “Rodrigues parameters” is much more subtle and is briefly discussed at the last section of the paper. A more in-depth discussion of the kinematics is left for future investigation.

The first part of the paper reviews the standard Cayley transform and it generalizes this transform to higher orders. There is no restriction on the dimension of the matrices involved, i.e., the results hold for $n \times n$ matrices. In the second part of the paper we apply these results to the case of interest to attitude dynamicists, i.e., the case $n = 3$.

Some notation and terminology is necessary in order to keep the discussion clear and terse. We use the standard mathematical notation $SO(n)$ to denote the space of proper orthogonal matrices of dimension $n \times n$. Invertible $n \times n$ matrices form the space $GL(n)$, the general linear group. The space of orthogonal matrices is denoted by $O(n)$ and it is the set of all (invertible) matrices $A \in GL(n)$ such that $A^T A = A A^T = I$. Clearly, if $A \in O(n)$ then

*Assistant Professor, Department of Mechanical, Aerospace and Nuclear Engineering. Member AIAA.

†Eppright Professor, Department of Aerospace Engineering. Fellow AIAA.

‡Graduate Student, Department of Aerospace Engineering. Student member AIAA.

$\det(A) = \pm 1$. The qualifier “proper” then refers to those orthogonal matrices with positive determinant, that is,

$$SO(n) = \{A \in Gl(n) : AA^T = I, \det(A) = +1\}$$

These matrices represent rotations, while the orthogonal matrices with determinant -1 represent reflections¹⁰. The space $SO(n)$ (as well as $Gl(n)$ and $O(n)$) forms a *group*. We will see later on that one can define a differential equation for elements of $SO(n)$. The solutions of this differential equation form trajectories (one-parameter subgroups) on $SO(n)$ and this differentiable structure makes $SO(n)$ actually a *Lie group* (i.e. a group with a differentiable manifold structure).

The space of $n \times n$ skew-symmetric matrices will be denoted by $so(n)$ That is,

$$so(n) = \{A \in \mathbb{R}^{n \times n} : A = -A^T\}$$

The space $so(n)$ is actually the tangent vector space to $SO(n)$ at the identity. This property can be easily verified by differentiating $A \in SO(n)$. Since $AA^T = I$ one has that

$$\frac{d}{dt}(AA^T) = 0 \Leftrightarrow \dot{A}A^T = -A\dot{A}^T$$

Evaluating the previous expression at $A = I$ one obtains that

$$\dot{A}\Big|_{A=I} = -\dot{A}^T\Big|_{A=I}$$

and so $\dot{A}\Big|_{A=I}$ is skew symmetric.

2. The Cayley Transform

Cayley’s transformation parameterizes a proper orthogonal matrix C as a function of a skew-symmetric matrix Q . It is, therefore, a map

$$\psi : so(n) \rightarrow SO(n) \quad (1)$$

The classical Cayley transform⁸ is given by

$$\begin{aligned} C = \psi(Q) &= (I - Q)(I + Q)^{-1} \\ &= (I + Q)^{-1}(I - Q) \end{aligned} \quad (2)$$

Since Q is skew-symmetric all its eigenvalues are pure imaginary. Thus, all the eigenvalues of the matrix $I + Q$ are nonzero and the inverse in Eq. (2) exists. The Cayley transform is therefore well-defined for all skew-symmetric matrices. The inverse transformation is identical and is given by

$$\begin{aligned} Q = \psi^{-1}(C) = \psi(C) &= (I - C)(I + C)^{-1} \\ &= (I + C)^{-1}(I - C) \end{aligned} \quad (3)$$

The inverse transformation is not defined when C has an eigenvalue at -1 , because in this case $\det(I + C) = 0$. Since C is orthogonal, all its eigenvalues lie on the unit circle

$$S^1 = \{(x_1, x_2) \in \mathbb{R}^2 : x_1^2 + x_2^2 = 1\} \quad (4)$$

Therefore $sp(C) \subset S^1$, where $sp(\cdot)$ denotes the spectrum of a matrix, and the transformation (3) requires that $-1 \notin sp(C)$. The same result is also shown in Ref. [7].

It is an easy exercise to show that C is orthogonal if Q is skew-symmetric. In order to show that the transformation (2) produces only proper orthogonal matrices, let us examine the determinant of C . Using Eq. (2) the determinant of C can be expressed as

$$\begin{aligned} \det(C) &= \det(I - Q)\det((I + Q)^{-1}) \\ &= \frac{\det(I - Q)}{\det(I + Q)} \end{aligned} \quad (5)$$

Since all the eigenvalues of Q are imaginary ($sp(Q) \subset \Im$) they are of the form $\pm i\lambda_j$. The spectral decomposition of the matrix Q then yields

$$Q = R^{-1}\Lambda R$$

where $\Lambda = \text{diag}(\pm i\lambda_j)$. (The matrix Q is normal and normal matrices are always diagonalizable¹¹.) Noting that $I \pm Q = R^{-1}(I \pm \Lambda)R$ we rewrite Eq. (5) as

$$\begin{aligned} \det(C) &= \frac{\det(R^{-1})\det(I - \Lambda)\det(R)}{\det(R^{-1})\det(I + \Lambda)\det(R)} = \frac{\det(I - \Lambda)}{\det(I + \Lambda)} \\ &= \frac{\prod_{j=1}^p (1 - i\lambda_j)(1 + i\lambda_j)}{\prod_{j=1}^p (1 + i\lambda_j)(1 - i\lambda_j)} \\ &= \frac{\prod_{j=1}^p (1 + \lambda_j^2)}{\prod_{j=1}^p (1 + \lambda_j^2)} = +1 \end{aligned}$$

where $2p$ is the number of nonzero (imaginary) eigenvalues of Q . Therefore $C \in SO(n)$ if $Q \in so(n)$ and thus, the Cayley transformation is injective (one-to-one) and surjective (onto) from the set of skew-symmetric matrices to the set of *proper* orthogonal matrices with no eigenvalue at -1 .

3. Cayley Transforms as Conformal Mappings

The three most important subsets of the complex numbers are the real numbers, the imaginary numbers, and the numbers with absolute value one (i.e., the numbers on the unit circle). Following the standard mathematical language, we use the symbols \mathbb{R} ,

$\Im = i\mathbb{R}$ and S^1 to denote these three sets, respectively. Trivially, these sets are subsets of the complex plane, denoted by \mathbb{C} . There is a very elegant analog between these three subsets of the complex plane and the $n \times n$ matrices⁹, i.e., the elements of $\mathbb{R}^{n \times n}$. This analog can be easily understood and appreciated as follows: An elementary result in matrix algebra states that every $n \times n$ matrix with real elements can be decomposed into the sum of a symmetric and a skew-symmetric matrix. For example, any $A \in \mathbb{R}^{n \times n}$ can be written as

$$A = \frac{A + A^T}{2} + \frac{A - A^T}{2} \quad (6)$$

It is easy to verify that the first matrix in Eq. (6) is symmetric and the second matrix is skew-symmetric. Symmetric matrices always have real eigenvalues and skew-symmetric matrices have always imaginary eigenvalues. Recall now that a complex number can always be decomposed into the sum of a real and an imaginary part. This parallelism between complex numbers and matrices allows one to treat the symmetric matrices as the “real numbers” and the skew-symmetric matrices as the “imaginary numbers” in the set of $\mathbb{R}^{n \times n}$ matrices⁹. In addition, recall that an orthogonal matrix in $\mathbb{R}^{n \times n}$ has all its eigenvalues on the unit circle. Drawing the previous parallelism even further we can therefore treat the orthogonal matrices as the “elements on the unit circle” in the space $\mathbb{R}^{n \times n}$. Similar statements can be made for the case of $n \times n$ matrices with complex entries (elements of $\mathbb{C}^{n \times n}$), where now hermitian, skew-hermitian and unitary matrices have to be used instead of symmetric, skew-symmetric and orthogonal matrices, respectively.

We intend to use this heuristic correspondence between complex numbers and $n \times n$ matrices in order to motivate and generalize the Cayley transform to higher order. Before we proceed, we briefly review some elements from complex function theory^{12,13}. First, recall that a (complex) function is analytic in an open set if it has a derivative at each point in that set. In particular, f is analytic at a point z_0 if it is analytic in a neighborhood of z_0 . Moreover, analytic functions have (uniformly) convergent power series expansions¹².

Definition 3.1 A transformation $w = f(z)$ where $w, z \in \mathbb{C}$ is said to be *conformal* at a point z_0 if f is analytic there and $f'(z_0) \neq 0$.

A conformal mapping is actually conformal at each point in a neighborhood of z_0 , since the analyticity of f at z_0 implies analyticity in a neighborhood of z_0 . Moreover, since f' is continuous at z_0 , it

follows that there is also a neighborhood of z_0 with $f'(z) \neq 0$ for all z in this neighborhood¹². It is a trivial consequence of the above definition that the composition of conformal mappings is also a conformal mapping.

A significant special class of conformal mappings in the complex plane is the class of *linear fractional transformations* (also called *bilinear transformations*) defined by

$$w = \frac{az + b}{cz + d}, \quad (ad - bc \neq 0) \quad (7)$$

An important property of the linear fractional transformations is that they always transform circles and lines into circles and lines¹². In this paper we are interested – in particular – in conformal transformations of the form (7) which map the unit circle on the imaginary axis and vice versa. One such transformation is given by $w = f(z)$ where

$$f(z) = \frac{1 - z}{1 + z} \quad (8)$$

It is an easy exercise to show that if $z \in \Im$ then $|w| = 1$, that is, $w \in S^1$ and thus, w is on the unit circle. Conversely, if $w \in S^1$ then the inverse transformation $z = f^{-1}(w)$ given by

$$f^{-1}(w) = \frac{1 - w}{1 + w} \quad (9)$$

implies that the real part of z is zero and thus, $z \in \Im$.

The inverse transformation (9) is defined everywhere except at $w = -1$. The point $w = -1$ is mapped to infinity (see Fig. 1). In fact, the map (8) introduces a one-to-one transformation $f : \Im \rightarrow S^1 \setminus \{-1\}$.




Figure 1: Bilinear transformation.

Let us now introduce the conformal mapping $g_n : S^1 \rightarrow S^1$ defined by

$$g_n(w) = w^n, \quad n = 2, 3, \dots \quad (10)$$

The function g_n is a mapping from the unit circle onto the unit circle. This transformation is only

locally injective. Therefore the inverse of g_n exists only locally. Given $\chi = e^{i\theta} \in S^1$ the solution of the equation

$$\chi = w^n, \quad n = 2, 3, \dots$$

yields that

$$w = e^{i\left(\frac{\theta+2k\pi}{n}\right)}, \quad k = 0, 1, 2, \dots, n-1 \quad (11)$$

Equation (11) shows that, in general, the equation $\chi = w^n$ has more than one solution. This result will turn out to be beneficial in section 5 when we discuss the application of higher order Cayley-transforms to attitude representations, because these roots can be used to avoid the inherent singularities of three-dimensional parameterizations of $SO(3)$. For $k = 0$ in Eq. (11) we get that $w = e^{i\frac{\theta}{n}}$. We will call this the *principal* n th root of χ .

The composition of the maps f and g_n is the function $h_n : \mathfrak{S} \rightarrow S^1$ defined by $h_n = g_n \circ f$, that is

$$h_n(z) = \left(\frac{1-z}{1+z}\right)^n \quad (12)$$

which maps the imaginary axis onto the unit circle. Similarly to g_n , this map is only locally invertible. A local inverse is obtained, for example, by setting $k = 0$ in Eq. (11), in which case we have that ($\chi = e^{i\theta}$)

$$z = e^{i\frac{\theta}{n}}$$

where

$$\theta = \arctan\left(i\frac{\bar{\chi} - \chi}{\bar{\chi} + \chi}\right)$$

and where bar denotes complex conjugate.

4. Higher Order Cayley Transforms

One of the most celebrated results in matrix algebra is the Cayley-Hamilton theorem. This theorem states that a matrix satisfies its own characteristic polynomial. An important consequence of this theorem is that, given any matrix $A \in \mathbb{R}^{n \times n}$ and an analytic function $F(z)$ inside a disk of radius r in the complex plane, one can unambiguously define the matrix-valued function $F(A)$ if the eigenvalues of A lie inside the disk of radius r . In other words, if F is given by

$$F(z) = \sum_{i=0}^{\infty} \alpha_i z^i, \quad |z| \leq r$$

then

$$F(A) = \sum_{i=0}^{\infty} \alpha_i A^i$$

and the previous series converges assuming that $|\lambda_j| \leq r$ where $\lambda_j \in sp(A)$ for $j = 1, 2, \dots, n$. Therefore, the matrix $F(A)$ is well-defined. Moreover, the eigenvalues of the matrix $F(A)$ are $F(\lambda_j)$ ($j = 1, 2, \dots, n$) (Ref. [11]).

Consider now the conformal mapping f from Eq. (8) which maps the imaginary axis on the unit circle. This function is analytic everywhere. According to the previous discussion, the matrix

$$f(Q) = (I - Q)(I + Q)^{-1} = (I + Q)^{-1}(I - Q) \quad (13)$$

is well-defined for $Q \in so(n)$ and, actually, $C = f(Q) \in SO(n)$. Comparison between the previous equation and Eq. (2) reveals that the Cayley transform can be viewed as a special case of a conformal mapping in the space of matrices.

We have seen that there is a natural correspondence between \mathfrak{S} and $so(n)$, as well as between S^1 and $SO(n)$. (We caution the the mathematically inclined reader to take these statements in the context of the discussion in section 3. We do not claim that this correspondence carries any more weight than providing one qualitative motivation for the generalization of certain complex analytic results to analogous results in the space of matrices). Following Eq. (12) we can also define a series of transformations $h_n : so(n) \rightarrow SO(n)$ by

$$h_n(Q) = (I - Q)^n (I + Q)^{-n} = (I + Q)^{-n} (I - Q)^n \quad (14)$$

where Q is a skew-symmetric matrix. It should be clear by now that $C = h_n(Q)$ is a proper orthogonal matrix, i.e., $C \in SO(n)$. We shall refer to the family of maps $h_n(Q)$ in Eq. (14) as *Higher Order Cayley Transforms*. The consequences of such a generalization in attitude representations will become apparent in the next section.

For now, let us concentrate on the inverse map $h_n^{-1} : SO(n) \rightarrow so(n)$. Since $h_n = g_n \circ f$ one obtains $h_n^{-1} = f^{-1} \circ g_n^{-1}$. The function f^{-1} is given by Eq. (9) which, when applied to a proper orthogonal matrix Q with no eigenvalue at -1 , gives the inverse of the classical (or first order) Cayley transform as in Eq. (3). The map $g_n^{-1} : SO(n) \rightarrow SO(n)$ on the other hand requires the n th root of an orthogonal matrix. First, we show that g_n^{-1} is well-defined in the sense that the n th root of a (proper) orthogonal matrix with no eigenvalue at -1 is also a (proper) orthogonal matrix with no eigenvalue at -1 . This will also prove that the composition of maps g_n^{-1} and f^{-1} is well-defined since the range of g_n^{-1} is in the domain of f^{-1} .

To this end, consider an orthogonal matrix $C \in SO(n)$ such that $\lambda \neq -1$ for all $\lambda \in sp(C)$. The

matrix C can be decomposed as follows

$$C = U\Theta U^* \quad (15)$$

for some unitary matrix U , where

$$\Theta = \text{blockdiag}(\Theta_1, \Theta_2, \dots, \Theta_{n-1}, +1) \quad (16)$$

if n is odd and

$$\Theta = \text{blockdiag}(\Theta_1, \Theta_2, \dots, \Theta_n) \quad (17)$$

if n is even, and

$$\Theta_j = \begin{bmatrix} e^{i\theta_j} & 0 \\ 0 & e^{-i\theta_j} \end{bmatrix}, \quad j = 1, \dots, n \quad (18)$$

The diagonal elements of the matrix Θ in Eq. (15) are the eigenvalues of C . The principal k th root of the matrix C is then given by

$$W = U\Theta_k U^* \quad (19)$$

where $W^k = C$ and

$$\Theta_k = \text{blockdiag}(\Theta_1^k, \Theta_2^k, \dots, \Theta_{n-1}^k, +1) \quad (20)$$

if n is odd and

$$\Theta_k = \text{blockdiag}(\Theta_1^k, \Theta_2^k, \dots, \Theta_n^k) \quad (21)$$

if n is even, and

$$\Theta_j^k = \begin{bmatrix} e^{i\frac{\theta_j}{k}} & 0 \\ 0 & e^{-i\frac{\theta_j}{k}} \end{bmatrix}, \quad j = 1, \dots, n \quad (22)$$

Since $e^{i\theta_j} \neq -1$ for all $j = 1, \dots, n$ ($n-1$) the angles $\theta_j \neq \pm 180$ deg and thus also $\frac{\theta_j}{k} \neq \pm 180$ deg for $k = 2, 3, \dots$ and thus $e^{i\frac{\theta_j}{k}} \neq -1$. Notice that in order to keep W proper we always choose the positive root of the eigenvalue $+1$.

5. Attitude Representations

In this section we concentrate on the ramifications of the previously developed results to attitude representations. Our motivation for investigating Cayley transforms in the first place, stems from the fact that proper orthogonal matrices represent rotations. In particular, $SO(3)$ is the configuration space of all three-dimensional rotations. In other words, every element of $SO(3)$ represents a physical rotation between two reference frames in \mathbb{R}^3 and conversely, every rotation can be represented by an element in $SO(3)$.

As a reference frame, viz. a body, rotates freely in the three-dimensional space, the corresponding

rotation matrix C traces a curve in $SO(3)$ such that $C(t) \in SO(3)$ for all $t \geq 0$. The differential equation characterizing this trajectory on $SO(3)$ is given by

$$\dot{C} = [\omega]C \quad (23)$$

where, given a vector $\omega = (\omega_1, \omega_2, \omega_3) \in \mathbb{R}^3$, the matrix $[\omega]$ is defined by

$$[\omega] = \begin{bmatrix} 0 & \omega_3 & -\omega_2 \\ -\omega_3 & 0 & \omega_1 \\ \omega_2 & -\omega_1 & 0 \end{bmatrix} \quad (24)$$

In the sequel we apply the results of the previous section in order to parameterize the rotation group. In particular, the series of conformal mappings from Eq. (14) provide a family of coordinates on $SO(3)$. Before undertaking this task we investigate another important conformal mapping.

5.1. The Exponential Map and the Euler Parameters

Linear fractional transformations are not the only class of conformal mappings from the imaginary axis onto the unit circle. The exponential map, defined by

$$w = \exp(z) = e^z \quad (25)$$

also maps \Im onto S^1 . Clearly, if $z = i\theta$ then $|z| = 1$. The inverse transformation is

$$z = \log w = i(\theta + 2n\pi), \quad n = 0, \pm 1, \pm 2, \dots$$

and is defined only locally.

We can therefore define the exponential map from the space of skew-symmetric matrices to the space of proper orthogonal matrices. This exponential map is defined, as usual, by

$$C = e^Q = \sum_{n=0}^{\infty} \frac{1}{n!} Q^n \quad (26)$$

and the series converges for every Q . For the three-dimensional case, the matrix $Q \in so(3)$ can be parameterized by

$$Q = \begin{bmatrix} 0 & \beta_3 & -\beta_2 \\ -\beta_3 & 0 & \beta_1 \\ \beta_2 & -\beta_1 & 0 \end{bmatrix} \quad (27)$$

As before, given a vector $\beta = (\beta_1, \beta_2, \beta_3) \in \mathbb{R}^3$ we will also use the notation $[\beta]$ to denote the skew-symmetric matrix in Eq. (27). Noticing that

$$[\beta]^2 = \beta\beta^T - \|\beta\|^2 I$$

one obtains that

$$[\beta]^{2k+1} = (-1)^k \|\beta\|^{2k} [\beta], \quad k = 0, 1, 2, \dots$$

and

$$[\beta]^{2k} = (-1)^k \|\beta\|^{2k} I - (-1)^k \|\beta\|^{2(k-1)} \beta \beta^T$$

Substituting the previous expressions in Eq. (26) we get Euler's formula³

$$C(\beta) = e^{[\beta]} = \cos \phi I + \sin \phi \frac{[\beta]}{\phi} + (1 - \cos \phi) \frac{\beta \beta^T}{\phi^2}$$

where $\phi = \|\beta\|$. Equivalently,

$$e^{[\beta]} = I + \sin \phi \frac{[\beta]}{\phi} + (1 - \cos \phi) \frac{[\beta]^2}{\phi^2} \quad (28)$$

Normalizing the vector β we get a unit vector

$$\hat{e} = \frac{\beta}{\|\beta\|}$$

or

$$\beta = \phi \hat{e} \quad (29)$$

Euler's theorem¹ states that any rotation can be represented by a finite rotation (principal rotation) about a single axis (principal axis). That is, the principal axis and the principal angle suffice to determine the rotation matrix. From a mathematical perspective this amounts to parameterizing every element in $SO(3)$ by the principal axis and the principal angle.

By letting the principal axis be along the direction of the unit vector \hat{e} and by letting the principal angle be ϕ as above, Eq. (28) shows how this parameterization is achieved. Clearly,

$$C(\phi, \hat{e}) = e^{\phi[\hat{e}]} \quad (30)$$

Moreover, introducing the *Euler parameter vector* $q = (q_0, q_1, q_2, q_3)$

$$q_0 = \cos \frac{\phi}{2}, \quad q_i = \hat{e}_i \sin \frac{\phi}{2}, \quad i = 1, 2, 3 \quad (31)$$

and substituting in Eq. (28) one obtains the well-known formula for the rotation matrix in terms of the Euler parameters

$$C(q) = \begin{bmatrix} q_0^2 + q_1^2 - q_2^2 - q_3^2 & 2(q_1 q_2 + q_0 q_3) \\ 2(q_1 q_2 - q_0 q_3) & q_0^2 - q_1^2 + q_2^2 - q_3^2 \\ 2(q_1 q_3 + q_0 q_2) & 2(q_2 q_3 - q_0 q_1) \\ & 2(q_1 q_3 - q_0 q_2) \\ & 2(q_2 q_3 + q_0 q_1) \\ & q_0^2 - q_1^2 - q_2^2 + q_3^2 \end{bmatrix} \quad (32)$$

Therefore, the Euler parameter representation is obtained by generalizing the conformal mapping in Eq. (25) to the space of matrices. Notice from Eq. (32) that $C(q) = C(-q)$ and both q and $-q$ can be used to describe the same physical orientation. This fact can be used to construct alternative, or "shadow", sets of kinematic parameters obtained via the Cayley transforms.

5.2. Rodrigues Parameters

Since the Euler parameters satisfy the additional constraint $q_0^2 + q_1^2 + q_2^2 + q_3^2 = 1$, one is naturally led to consider the elimination of this constraint, thus reducing the number of coordinates from four to three. The Rodrigues parameters achieve this by defining

$$\rho_j = \frac{q_j}{q_0}, \quad j = 1, 2, 3 \quad (33)$$

The three parameters ρ_1, ρ_2, ρ_3 then provide a three-dimensional parameterization of $SO(3)$. The inverse transformation of Eq. (33) is given by

$$q_0 = \frac{1}{(1 + \hat{\rho}^2)^{\frac{1}{2}}}, \quad q_j = \frac{\rho_j}{(1 + \hat{\rho}^2)^{\frac{1}{2}}}, \quad j = 1, 2, 3 \quad (34)$$

where $\hat{\rho}^2 = \rho_1^2 + \rho_2^2 + \rho_3^2$. The Rodrigues parameters are related to the principal axis and angle through the equation

$$\rho = \tan \frac{\phi}{2} \hat{e}$$

The rotation matrix in terms of the Rodrigues parameters can be easily computed using Eq. (32) and Eq. (34).

$$C(\rho) = \frac{1}{1 + \hat{\rho}^2} \begin{bmatrix} 1 - \hat{\rho}^2 + 2\rho_1^2 & 2(\rho_1 \rho_2 + \rho_3) \\ 2(\rho_1 \rho_2 - \rho_3) & 1 - \hat{\rho}^2 + 2\rho_2^2 \\ 2(\rho_3 \rho_1 + \rho_2) & 2(\rho_2 \rho_3 - \rho_1) \\ & 2(\rho_3 \rho_1 - \rho_2) \\ & 2(\rho_2 \rho_3 + \rho_1) \\ & 1 - \hat{\rho}^2 + 2\rho_3^2 \end{bmatrix} \quad (35)$$

It is remarkable the fact that the previous parameterization of $SO(3)$ can also be achieved by means of the Cayley transformation in Eq. (2). Indeed, if we introduce the skew-symmetric matrix

$$R = -[\rho] = \begin{bmatrix} 0 & -\rho_3 & \rho_2 \\ \rho_3 & 0 & -\rho_1 \\ -\rho_2 & \rho_1 & 0 \end{bmatrix}$$

the transformation

$$C = (I - R)(I + R)^{-1} = (I + R)^{-1}(I - R) \quad (36)$$

produces exactly the matrix in Eq. (35). Therefore the classical Cayley-Rodrigues parameters representation is obtained by generalizing the conformal mapping in Eq. (8) to the space of matrices.

5.3. Modified Rodrigues Parameters

The normalization in Eq. (33) is not the only possible one. A more judicious normalization for eliminating the Euler parameter constraint is through stereographic projection^{12,13,14}. Using this approach, the new variables

$$\sigma_j := \frac{q_j}{1 + q_0}, \quad j = 1, 2, 3 \quad (37)$$

provide coordinates on $SO(3)$. These parameters are referred to in the literature as the Modified Rodrigues parameters³ and have distinct advantages over the classical Rodrigues parameters. In particular, while the Rodrigues parameters do not allow eigenaxis rotations of more than 180 deg, the Modified Rodrigues parameters allow for eigenaxis rotations of upto 360 deg^{6,7,14,15,16}. This can be immediately deduced by the corresponding relationship between σ and the principal axis and angle

$$\sigma = \tan \frac{\phi}{4} \hat{e}$$

which is well-behaved for $0 \leq \phi < 2\pi$. Since both q and $-q$ describe the same physical orientation (recall the discussion at the end of section 5.1), a second set of parameters defined by

$$\sigma_j^s := -\frac{q_j}{1 - q_0}, \quad j = 1, 2, 3$$

referred to as the ‘‘shadow’’ set¹⁵, can be used to describe the same physical orientation. These parameters are also given by

$$\sigma^s = -\frac{1}{\tan \frac{\phi}{4}} \hat{e}$$

The transformation between σ and σ^s is given by¹⁵

$$\sigma^s = -\frac{\sigma}{\hat{\sigma}^2} \quad (38)$$

where $\hat{\sigma}^2 = \sigma^T \sigma = \sigma_1^2 + \sigma_2^2 + \sigma_3^2 = \tan^2 \frac{\phi}{4}$.

The rotation matrix associated with the Modified Rodrigues Parameters is given by

$$C(\sigma) = \frac{1}{1 + \hat{\sigma}^2} \begin{bmatrix} 4\Sigma_1 + \tilde{\Sigma}^2 & 8\sigma_1\sigma_2 + 4\sigma_3\tilde{\Sigma} & & \\ 8\sigma_1\sigma_2 - 4\sigma_3\tilde{\Sigma} & 4\Sigma_2 + \tilde{\Sigma}^2 & & \\ 8\sigma_1\sigma_3 + 4\sigma_2\tilde{\Sigma} & 8\sigma_2\sigma_3 - 4\sigma_1\tilde{\Sigma} & & \\ & 8\sigma_1\sigma_3 - 4\sigma_2\tilde{\Sigma} & & \\ & 8\sigma_2\sigma_3 + 4\sigma_1\tilde{\Sigma} & & \\ & 4\Sigma_3 + \tilde{\Sigma}^2 & & \end{bmatrix} \quad (39)$$

where $\tilde{\Sigma} = 1 - \hat{\sigma}^2$ and $\Sigma_j = -\hat{\sigma}^2 + 2\sigma_j^2$, $j = 1, 2, 3$.

In Ref. [7] it was shown that these parameters are defined by a Cayley transformation of second order. That is, if

$$S = -[\sigma] = \begin{bmatrix} 0 & -\sigma_3 & \sigma_2 \\ \sigma_3 & 0 & -\sigma_1 \\ -\sigma_2 & \sigma_1 & 0 \end{bmatrix} \quad (40)$$

then the transformation

$$C = (I - S)^2(I + S)^{-2} = (I + S)^{-2}(I - S)^2 \quad (41)$$

produces exactly the matrix in Eq. (39). Notice that the inverse of the transformation (41) is not unique and it requires the square root of an orthogonal matrix. Given $C \in SO(3)$ we find a matrix W such that

$$C = W^2 \quad (42)$$

Once a matrix W is calculated, the skew-symmetric matrix S containing the Modified Rodrigues parameters is computed from

$$S = (I - W)(I + W)^{-1} = (I + W)^{-1}(I - W) \quad (43)$$

Reference [7] outlines this approach. To every orthogonal matrix corresponds a principal angle and a principal direction according to Eq. (30). From Eqs. (30) and (42) one therefore has that

$$W = e^{\frac{\phi}{2}[\hat{e}]} \quad (44)$$

and W has half the principal angle of C . It should be apparent now how the Modified Rodrigues parameters double the domain of validity of the parameterization by taking the square of the classical Cayley transform.

This observation motivates the search of higher dimensional Cayley transforms for attitude representations. Such transformations are expected to increase the domain of validity even further. This is the topic of the next section.

5.4. Higher Order Rodrigues Parameters

According to the discussion in the previous section one expects that higher order Cayley transformations will increase the domain of validity of the corresponding parameters. The main task of this section is to derive these higher order parameters and find their connections to the Rodrigues parameters, the Modified parameters and the Euler parameters. To this end, consider first the fourth order Cayley transform defined by

$$C = (I - T)^4(I + T)^{-4} \quad (45)$$

for some skew-symmetric matrix

$$T = -[\tau] = \begin{bmatrix} 0 & -\tau_3 & \tau_2 \\ \tau_3 & 0 & -\tau_1 \\ -\tau_2 & \tau_1 & 0 \end{bmatrix} \quad (46)$$

We know that the matrix C is (proper) orthogonal.

Recall from the results of section 3 that if F is analytic function, then the eigenvalues of the matrix $F(A)$ are given by $F(\lambda_j)$ where λ_j are the eigenvalues of A . It is an easy exercise to show that the eigenvalues of the skew-symmetric matrix in Eq. (46) are given by

$$0, \quad \pm i(\tau_1^2 + \tau_2^2 + \tau_3^2)^{\frac{1}{2}} \quad (47)$$

Similarly, the eigenvalues of the matrix S in Eq. (40) are given by

$$0, \quad \pm i(\sigma_1^2 + \sigma_2^2 + \sigma_3^2)^{\frac{1}{2}} \quad (48)$$

Let λ_τ denote an eigenvalue of T and λ_σ an eigenvalue of S . Comparing Eqs. (41) and (45) one sees that the matrices S and T are related by

$$(I - S)(I + S)^{-1} = (I - T)^2(I + T)^{-2} \quad (49)$$

This suggests that λ_σ and λ_τ are related by

$$\frac{1 - \lambda_\sigma}{1 + \lambda_\sigma} = \left(\frac{1 - \lambda_\tau}{1 + \lambda_\tau} \right)^2 \quad (50)$$

or

$$1 + \lambda_\sigma = \frac{(1 + \lambda_\tau)^2}{1 + \lambda_\tau^2}$$

Solving for λ_σ one obtains that

$$\lambda_\sigma = \frac{2\lambda_\tau}{1 + \lambda_\tau^2}$$

Substituting the expressions for λ_σ and λ_τ from Eqs. (47) and (48) in the previous equation one obtains that

$$\pm i(\sigma_1^2 + \sigma_2^2 + \sigma_3^2)^{\frac{1}{2}} = 2 \frac{\pm i(\tau_1^2 + \tau_2^2 + \tau_3^2)^{\frac{1}{2}}}{1 - \tau_1^2 - \tau_2^2 - \tau_3^2}$$

Upon squaring this expression one obtains

$$\sigma_1^2 + \sigma_2^2 + \sigma_3^2 = 4 \frac{\tau_1^2 + \tau_2^2 + \tau_3^2}{(1 - \tau_1^2 - \tau_2^2 - \tau_3^2)^2}$$

This equation suggests that σ and τ are related by

$$\sigma_j = \pm \frac{2\tau_j}{1 - \tau_1^2 - \tau_2^2 - \tau_3^2}, \quad j = 1, 2, 3 \quad (51)$$

Arbitrarily, and without loss of generality, we choose the solution with the plus sign. Substitution in S

and computing C from Eq. (41) verifies the expression in Eq. (51).

The relation between τ and q is obtained by observing that

$$\frac{2\tau_j}{1 - \tau_1^2 - \tau_2^2 - \tau_3^2} = \frac{q_j}{1 + q_0}, \quad j = 1, 2, 3 \quad (52)$$

Using the shorthand notation $\hat{\tau}^2 = \tau_1^2 + \tau_2^2 + \tau_3^2$ the previous expression can be written as

$$\frac{2\tau_j}{1 - \hat{\tau}^2} = \frac{q_j}{1 + q_0}, \quad j = 1, 2, 3$$

Therefore,

$$4 \frac{\hat{\tau}^2}{(1 - \hat{\tau}^2)^2} = \frac{q_1^2 + q_2^2 + q_3^2}{(1 + q_0)^2}$$

or

$$\begin{aligned} \left(\frac{1 + \hat{\tau}^2}{1 - \hat{\tau}^2} \right)^2 &= \frac{q_1^2 + q_2^2 + q_3^2 + (1 + q_0)^2}{(1 + q_0)^2} \\ &= \frac{2(1 + q_0)}{(1 + q_0)^2} = \frac{2}{1 + q_0} \end{aligned} \quad (53)$$

or that

$$\frac{1 + \hat{\tau}^2}{1 - \hat{\tau}^2} = \pm \frac{\sqrt{2}}{\sqrt{1 + q_0}}$$

and thus,

$$\frac{2}{1 - \hat{\tau}^2} = \frac{\pm\sqrt{2} + \sqrt{1 + q_0}}{\sqrt{1 + q_0}}$$

Using now Eq. (52) one finally obtains that

$$\tau_j = \frac{q_j}{1 + q_0 \pm \sqrt{2(1 + q_0)}}, \quad j = 1, 2, 3 \quad (54)$$

Conversely, from Eq. (53) one obtains that

$$1 + q_0 = 2 \left(\frac{1 - \hat{\tau}^2}{1 + \hat{\tau}^2} \right)^2 \quad (55)$$

and using Eq. (52) that

$$q_j = \frac{4\tau_j(1 - \hat{\tau}^2)}{(1 + \hat{\tau}^2)^2}, \quad j = 1, 2, 3$$

From Eq. (55) we also have that

$$q_0 = 2 \left(\frac{1 - \hat{\tau}^2}{1 + \hat{\tau}^2} \right)^2 - 1 = \frac{(1 - 6\hat{\tau}^2 + \hat{\tau}^4)}{(1 + \hat{\tau}^2)^2}$$

where $\hat{\tau}^4 = (\hat{\tau}^2)^2$. Letting $W = (I - T)(I + T)^{-1}$ and since $C = W^4$ one obtains that

$$W = e^{\frac{\phi}{4}[\hat{\epsilon}]}$$

where ϕ is the principal angle of C . Moreover, using the definition of the Euler parameters from Eq. (31) one obtains the following result for the τ parameters

$$\tau = \frac{\sin \frac{\phi}{2}}{1 + \cos \frac{\phi}{2} \pm \sqrt{2(1 + \cos \frac{\phi}{2})}} \hat{e} \quad (56)$$

where \hat{e} is the unit vector along the principal axis. Using the trigonometric identity $\cos \frac{\phi}{2} = 2 \cos^2 \frac{\phi}{4} - 1$, the previous equation reduces to

$$\tau = \frac{\sin \frac{\phi}{2}}{1 + \cos \frac{\phi}{2} \pm 2 \cos \frac{\phi}{4}} \hat{e} \quad (57)$$

Keeping the plus sign, Eq. (57) can be further reduced to the simple formula

$$\tau_+ = \tan \frac{\phi}{8} \hat{e}, \quad (-4\pi < \phi < 4\pi) \quad (58)$$

From Eq. (58) it is apparent that τ is proportional to the principal rotation axis, like the classical and the Modified Rodrigues parameters, where now the proportionality factor is $f(\phi) = \tan \frac{\phi}{8}$. A plot of $f(\phi)$ is shown in Fig. 2.




Figure 2: Plot of $f(\phi)$.

Equation (58) is reassuring, since it proves that the τ parameters indeed behave as “higher order” Rodrigues parameters which can be used to “linearize” the domain of validity of the kinematic parameterization. By this, we mean that Eq. (58) behaves almost linearly as a function of the principal angle ϕ (especially in the region $-\pi/8 \leq \phi \leq \pi/8$); see also Fig. 3.

If we choose the minus sign in Eq. (56) we obtain that

$$\tau_- = -\frac{1}{\tan \frac{\phi}{8}} \hat{e}, \quad (0 < \phi < 8\pi) \quad (59)$$

Moreover, reversing the signs of the Euler parameters in Eq. (54), one obtains that the τ parameters

have a unique set of “shadow” parameters like the Modified Rodrigues parameters¹⁵. These parameters are obtained by setting

$$\tau^s = \frac{-\sin \frac{\phi}{2}}{1 - \cos \frac{\phi}{2} \pm 2 \sin \frac{\phi}{4}} \hat{e} \quad (60)$$

It can be easily verified that the corresponding “shadow” parameters reduce to

$$\tau_+^s = \frac{\tan \frac{\phi}{8} - 1}{\tan \frac{\phi}{8} + 1} \hat{e} \quad (-2\pi < \phi < 6\pi) \quad (61)$$

and

$$\tau_-^s = \frac{1 + \tan \frac{\phi}{8}}{1 - \tan \frac{\phi}{8}} \hat{e} \quad (-6\pi < \phi < 2\pi) \quad (62)$$

As the original τ parameters approach $+1$, the associated “shadow” parameters τ^s approach zero and vice versa. The general transformation between the original and the “shadow” set is given by

$$\tau^s = -\tau \left(\frac{1 - \hat{\tau}^2}{2\hat{\tau}^2 + (1 + \hat{\tau}^2)\hat{\tau}} \right) \quad (63)$$

where $\hat{\tau} = (\tau^2)^{\frac{1}{2}}$. Equations (58), (59), (61) and (62) can be used in order to compute the four distinct roots of Eq. (45). Note also that Eqs. (58), (61), (59) and (62) can be also written in the form

$$\tau = \tan \left(\frac{\phi}{8} - k \frac{\pi}{4} \right) \hat{e}, \quad k = 0, 1, 2, 3$$

respectively.

The “shadow” parameter set τ^s is shown side-by-side with the original τ parameters in Fig. 3. The shadow set is plotted in grey color. Figure 3 also shows that τ parameters are indeed very linear for small rotations within ± 180 deg.

As with the Modified Rodrigues parameters (and other stereographic parameters¹⁵), these “shadow” parameters represent the same physical orientation as the original set and abide by the same differential kinematic equation. They could be used to avoid the problems of approaching the ± 720 deg principal rotation. By switching to the shadow trajectory, all numerical problems would be avoided. Having, however, a principal rotation range of ± 720 deg is really more than needed. Limiting the principal rotations to be within ± 180 deg would suffice and be much more attractive. As the magnitude of τ approaches $\tan \frac{\pi}{8}$ then one would simply switch the τ to their “shadow” set. Having $|\tau| = \tan \frac{\pi}{8}$ corresponds to $q_0 = 0$. From Eq. (54) one can then see




Figure 3: Comparison of original and “shadow” τ parameters.

that at this point, the two sets of parameters are related by $\tau = -\tau^s$. The combined set of original and “shadow” τ parameters would provide a set of attitude coordinates which are “very linear” with respect to the principal rotation angle, more so even than the Modified Rodrigues parameters. We note in passing that the previous approach can be easily extended to any Cayley transform of order 2^k , since Eqs. (49) and (50) can be used iteratively.

For the third order Cayley transform we have that

$$C = (I - P)^3(I + P)^{-3} = (I + P)^{-3}(I - P)^3 \quad (64)$$

where $P = -[p]$ and $p = (p_1, p_2, p_3)$ the corresponding parameters. If λ_ρ and λ_p denote the respective eigenvalues of the skew-symmetric matrices R and P then, using Eqs. (36) and (64), they must be related by

$$\frac{1 - \lambda_\rho}{1 + \lambda_\rho} = \left(\frac{1 - \lambda_p}{1 + \lambda_p} \right)^3$$

or, upon expanding the previous equality

$$\frac{1 - \lambda_\rho}{1 + \lambda_\rho} = \frac{1 - \lambda_p^3 + 3\lambda_p^2 - 3\lambda_p}{1 + \lambda_p^3 + 3\lambda_p^2 + 3\lambda_p}$$

thus

$$1 + \lambda_\rho = \frac{(1 + \lambda_p)^3}{1 + 3\lambda_p^2}$$

Solving for λ_ρ we obtain

$$\lambda_\rho = \frac{\lambda_p(3 + \lambda_p^2)}{1 + 3\lambda_p^2}$$

The previous equation suggests that ρ_j and p_j are related by

$$\rho_j = \pm \frac{p_j(3 - p_1^2 - p_2^2 - p_3^2)}{1 - 3(p_1^2 + p_2^2 + p_3^2)}, \quad j = 1, 2, 3$$

In order to get the relation of p to the Euler parameter vector one can set

$$\frac{p_j(3 - p_1^2 - p_2^2 - p_3^2)}{1 - 3(p_1^2 + p_2^2 + p_3^2)} = \frac{q_j}{q_0} \quad (65)$$

and solve for $\hat{p}^2 = p_1^2 + p_2^2 + p_3^2$. After some algebraic calculations, it is not difficult to show that, in fact,

$$\frac{(\hat{p}^2 + 1)^3}{(1 - 3\hat{p}^2)^2} = \frac{1}{q_0^2} \quad (66)$$

Solution of the previous equation for \hat{p}^2 requires the solution of a cubic equation. Once \hat{p}^2 is known however, it can be substituted into Eq. (65) to get the desired result. Actually, from Eqs. (65) and (66) we have that

$$q_0 = \frac{1 - 3\hat{p}^2}{(1 + \hat{p}^2)^{\frac{3}{2}}}, \quad q_j = \pm \frac{p_j(3 - \hat{p}^2)}{(1 + \hat{p}^2)^{\frac{3}{2}}}, \quad j = 1, 2, 3$$

Letting $W = (I - P)(I + P)^{-1}$ then since $C = W^3$ one obtains that

$$W = e^{\frac{\phi}{3}[\hat{e}]}$$

where ϕ is the principal angle of C .

6. Kinematics

The kinematic equations in terms of the τ parameters can be computed as follows. From Eqs. (23) and (45) we have that

$$\begin{aligned} \dot{C} &= \frac{d}{dt}[(I - T)^4](I + T)^{-4} + (I - T)^4 \frac{d}{dt}[(I + T)^{-4}] \\ &= S(\omega)(I - T)^4(I + T)^{-4} \end{aligned}$$

or that

$$\frac{d}{dt}[(I - T)^4] - C(T) \frac{d}{dt}[(I + T)^4] = S(\omega)(I - T)^4 \quad (67)$$

where we have used the fact that

$$\frac{d}{dt}A^{-1} = -A^{-1} \left(\frac{d}{dt}A \right) A^{-1}$$

for any square matrix A . Using also the fact that

$$\frac{d}{dt}A^n = \sum_{j=0}^{n-1} A^j \left(\frac{d}{dt}A \right) A^{n-j-1}$$

and performing the differentiations in the left-hand-side of Eq. (67), one obtains a set of nine linear equations in terms of $\dot{\tau}_1, \dot{\tau}_2,$ and $\dot{\tau}_3$. Similarly, the right-hand-side of Eq. (67) is linear in terms of $\omega_1, \omega_2, \omega_3$.

Choosing three (independent) equations out of these nine, we get a linear system of the form

$$V(\tau)\dot{\tau} = U(\tau)\omega$$

Solving for $\dot{\tau}$ we finally get that the kinematic equations for the τ orientation parameters are given by

$$\frac{d\tau}{dt} = V^{-1}(\tau)U(\tau)\omega = G(\tau)\omega$$

where the matrix $G(\tau)$ is given by

$$G(\tau) = \frac{1}{1 - \hat{\tau}^2} \begin{bmatrix} T_1 + \tau_3^2 \tau_2^2 - 3(\tau_2^2 + \tau_3^2) \\ 2\tau_3(1 - \hat{\tau}^2) + \tau_1 \tau_2(3 - \hat{\tau}^2) \\ -2\tau_2(1 - \hat{\tau}^2) + \tau_1 \tau_3(3 - \hat{\tau}^2) \\ -2\tau_3(1 - \hat{\tau}^2) + \tau_1 \tau_2(3 - \hat{\tau}^2) \\ T_2 + \tau_1^2 \tau_3^2 - 3(\tau_3^2 + \tau_1^2) \\ 2\tau_1(1 - \hat{\tau}^2) + \tau_2 \tau_3(3 - \hat{\tau}^2) \\ 2\tau(1 - \hat{\tau}^2) + \tau_1 \tau_3(3 - \hat{\tau}^2) \\ -2\tau_1(1 - \hat{\tau}^2) + \tau_2 \tau_3(3 - \hat{\tau}^2) \\ T_3 + \tau_1^2 \tau_2^2 - 3(\tau_1^2 + \tau_2^2) \end{bmatrix} \quad (68)$$

and $T_j = \frac{1}{2}(1 + \tau_1^4 + \tau_2^4 + \tau_3^4 - 2\tau_j^4)$, $j = 1, 2, 3$. This equation can be written more compactly in a vector form as follows

$$\frac{d\tau}{dt} = \frac{1}{8(1 - \hat{\tau}^2)} [2(3 - \hat{\tau}^2)\tau\tau^T - 4(1 - \hat{\tau}^2)[\tau] + (1 - 6\hat{\tau}^2 + \hat{\tau}^4)I] \omega \quad (69)$$

These kinematic equations are not as simple as the corresponding kinematic equations for the Rodrigues or the Modified Rodrigues parameters^{7,14}. Moreover, there is an apparent singularity at $\hat{\tau} = \pm 1$, equivalently at $\phi = \pm 2\pi$. The limiting behavior of these equations as $\hat{\tau} \rightarrow \pm 1$ will be determined through further analytical and numerical studies. At any rate, because of the near-linear behavior between ϕ and the magnitude of τ as seen in Fig. 2, for small principal angles, Eq. (69) is expected to behave in a more “linear-like” fashion than either the Cayley-Rodrigues or the Modified Rodrigues parameters.

Similarly, for the third order Cayley parameters, one can derive the following kinematic equations

$$\frac{dp}{dt} = \frac{1}{6(3 - \hat{p}^2)} [(11 - \hat{p}^2)pp^T - 3(3 - \hat{p}^2)[p] + 3(1 - 3\hat{p}^2)I] \omega \quad (70)$$

These equations can be derived starting from Eqs. (23) and (64) and using similar arguments as before. Singularities for the p parameters are encountered at $\hat{p} = \pm\sqrt{3}$. As before, further analysis is required to determine the limiting behavior of this system as $\hat{p} \rightarrow \pm\sqrt{3}$.

7. Numerical Example

In order to demonstrate the potential benefits or drawbacks of the previous kinematic parameters the following simulation was performed. We integrated Eqs. (69) as well as the corresponding kinematic equations in terms of the Cayley-Rodrigues (ρ) and the Modified Rodrigues parameters (σ) starting from the zero orientation and subject to the constant angular velocity vector $\omega = (0.25, 0.4, -0.1)$ (rad/sec). This corresponds to a linearly increasing value of the principal angle ϕ . The results of the simulations are shown in Fig. 4. This figure actually shows only the first components of the kinematic parameter vectors, as the other two components exhibit similar behavior.




Figure 4: Orientation parameter comparison.

As it is evident from this figure, the classical and the Modified Rodrigues parameters encounter the singularity earlier than the τ parameters. We note, however, that since discontinuities in the parameter description are typically acceptable in applications, the Modified Rodrigues parameters can be made to avoid the singularity altogether by simply switching to their “shadow” set¹⁵. The same also holds for the τ parameters via Eq. (63). Figure 5 shows the simulation where the parameters σ and τ are allowed to switch to their respective “shadow” sets. Although the points of switching are arbitrary and can be chosen according to the particular application, a reasonable choice is to switch when the parameters and the corresponding “shadow” set have opposite signs. This will ensure continuity of the magnitude. From Eqs. (38) and (63) this occurs when $\phi = k\pi$, $k = \pm 1, \pm 2, \dots$. This is the situation depicted in Fig. 5. The τ parameters are shown in solid line, and the σ parameters are shown in dashed line.




Figure 5: Orientation parameter and their “shadow” sets.

Since the classical Rodrigues parameters do not have an associated “shadow” set (better, the shadow set coincides with the original parameters), only the σ and τ parameters are plotted in Fig. 5.

8. Conclusions

We have extended the classical Cayley transform which maps skew-symmetric matrices to proper orthogonal matrices to higher orders. The approach is based on the observation that Cayley transforms can be viewed as generalized conformal (bilinear) mappings in the space of matrices. The Euler parameters, the Rodrigues parameters and the Modified Rodrigues parameters follow as special cases of this approach. In addition, we generate a family of higher order “Rodrigues parameters” which could be used as coordinates for the rotation group. It still remains, however, to determine the applicability of these higher order parameters in realistic attitude problems.

9. References

- [1] Hughes, P. C., *Spacecraft Attitude Dynamics*. New York, John Wiley & Sons, 1986.
- [2] Wertz, J. R., *Spacecraft Attitude Determination and Control*. Dordrecht, Holland, D. Reidel Publishing Company, 1980.
- [3] Shuster, M. D., “A Survey of Attitude Representations,” *Journal of the Astronautical Sciences*, Vol. 41, No. 4, 1993, pp. 439–517.
- [4] Special Issue on Attitude Representations, *Journal of the Astronautical Sciences*, Vol. 41, No. 4, 1993.
- [5] Wiener, T. F., *Theoretical Analysis of Gimbal-less Inertial Reference Equipment Using Delta-Modulated Instruments*. PhD thesis, Massachusetts Institute of Technology, Cambridge, Massachusetts, March 1962.
- [6] Marandi, S. R. and Modi, V., “A Preferred Coordinate System and the Associated Orientation Representation in Attitude Dynamics,” *Acta Astronautica*, Vol. 15, No. 11, 1987, pp. 833–843.
- [7] Schaub, H., Tsiotras, P., and Junkins, J. L., “Principal Rotation Representations of Proper $N \times N$ Orthogonal Matrices,” *International Journal of Engineering Science*, Vol. 33, No. 15, 1995, pp. 2277–2295.
- [8] Junkins, J. L. and Kim, Y., *Dynamics and Control of Flexible Structures*. New York, AIAA, 1993.
- [9] Halmos, P. R., *Finite Dimensional Vector Spaces*, Vol. 7 of *Annals of Mathematics Studies*. Princeton, NJ, Princeton University Press, 1953.
- [10] Curtis, M. L., *Matrix Groups*. New York, Springer-Verlag, 1979.
- [11] Horn, R. and Johnson, C. R., *Matrix Analysis*. Cambridge, United Kingdom, Cambridge University Press, 1985.
- [12] Churchill, R. V. and Brown, J. W., *Complex Variables and Applications*. New York, McGraw Hill, 1990.
- [13] Conway, J. B., *Functions of One Complex Variable*. New York, Springer Verlag, 1978.
- [14] Tsiotras, P., “On New Parameterizations of the Rotation Group in Attitude Kinematics,” Technical Report, Dept. of Aeronautics & Astronautics, Purdue University, West Lafayette, IN, January 1994.
- [15] Schaub, H. and Junkins, J. L., “Stereographic Orientation Parameters for Attitude Dynamics: A Generalization of the Rodrigues Parameters,” in *AAS/AIAA Space Flight Mechanics Conference*, Feb. 13–15, 1995. Albuquerque, NM.
- [16] Tsiotras, P., “New Control Laws for the Attitude Stabilization of Rigid Bodies,” in *13th IFAC Symposium on Automatic Control in Aerospace*, pp. 316–321, Sept. 14–17, 1994. Palo Alto, CA.